

AD-A056 492

OFFICE OF THE SURGEON GENERAL (ARMY) WASHINGTON D C
TOXICITY AND TREATMENT OF SEA SNAKE ENVENOMATION, (U)
JUN 78 J A VICK, G V PICKWELL


F/G 6/5

UNCLASSIFIED

NL

1 of 1
AD
A056 492





LEVEL II

VICK & PICKWELL

AD A 056492

TOXICITY AND TREATMENT OF SEA SNAKE ENVENOMATION (U)

JAMES A. VICK, LTC MSC
GEORGE V. PICKWELL Ph D.
OFFICE OF THE SURGEON GENERAL
WASHINGTON, D.C. 20310

DDC
RECEIVED
JUL 12 1978
D

11 JUN 1978

INTRODUCTION.

In recent years, increasing interest has been directed to the sea snakes of the family Hydrophidae with special attention to their behavior, general biology, and the chemical nature of their venoms. Several television features concerning sea snakes, together with newspaper and magazine articles, have generated substantial public interest in these animals (1-3).

Perhaps the greatest current interest in sea snakes lies in the probability of one species, *Pelamis platurus* (the yellow bellied sea snake), being an early immigrant into the Caribbean Sea from the Pacific Ocean when a Middle American sea-level canal is constructed (4-6).

At present no resident, reproducing population of sea snakes of any species has been described from the Caribbean Sea or the Atlantic Ocean.

Concern on the part of members of the scientific community stems from the lack of adequate knowledge with which to judge the probable effect that *Pelamis*, a fish-eating snake, would have on Caribbean pelagic ecology. A second and more immediate problem is the threat to human safety, particularly in regions frequented by tourists. Some scientists have predicted disastrous economic consequences to Caribbean tourist centers (presumed reduction in tourist trade) arising from the presence of this sea snake (6,7). The problem in this latter point of concern seemed to us to center about the following points: (1) the actual potency of this sea snake's venom; (2) the quantity of venom available for delivery by individual snakes; (3) the degree of aggressiveness or willingness to bite displayed by

78 06 09 090

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

266 700

VICK & PICKWELL

this snake, as well as the total number of snakes; and (4) the degree of fear expressed by people who encounter this snake. In attempting to answer these questions, we hope to shed some light on the true seriousness of the threat to man posed by marine snakes in a new habitat. This paper is a report on our findings, and includes an evaluation of the currently fielded sea snake antivenin

MATERIALS AND METHODS.

A. Sea Snakes.

Our specimens of *Pelamis platurus* were all collected within the confines of Bahia Banderas, Mexico. This is a deep water bay indenting the Pacific coast some 30 miles and having a mouth 10 to 12 miles across. The town of Puerto Vallarta at the foot of the bay (population approximately 25,000) is a tourist resort popular among Mexican and American citizens alike.

The distribution of sea snakes in this area is similar to that which might be expected in the area around the proposed sea way canal.

While our research vessels were hove-to or at anchor during the Naval Undersea Center cruises "MINOXI (July 1970) PELACAN I (March 1971), and MINOXII (March 1972)", we collected the sea snakes by dip-netting. Captured specimens were held aboard ship in gimbaled aquaria supplied with air bubblers and filters. The snakes were usually milked of their venom and tagged within moments of capture, but some were held for several days before their first milking

When our holding capacity for live snakes aboard ship were reached (30 to 40 adult specimens), we continued milking newly captured snakes, immediately returning them to the water. Upon the ship's return to San Diego, the captured snakes were placed in aquaria supplied with seawater and filters; aquaria were located in a temperature-controlled room at 80 degrees F. The snakes were fed live goldfish at intervals of 7 to 10 days and were milked every 3 to 4 weeks.

B. Obtaining Venom.

The adult of the sea snake has fangs only 2 mm in length. We therefore followed the method of Barne⁽⁸⁾, using capillary tubes of known volume for obtaining the venom. The milking operation involved three operators one restraining and manipulating the snake while the

ACCESSION NO.	2718	2719	2720	2721	2722	2723	2724	2725	2726	2727	2728	2729	2730	2731	2732	2733	2734	2735	2736	2737	2738	2739	2740	2741	2742	2743	2744	2745	2746	2747	2748	2749	2750	2751	2752	2753	2754	2755	2756	2757	2758	2759	2760	2761	2762	2763	2764	2765	2766	2767	2768	2769	2770	2771	2772	2773	2774	2775	2776	2777	2778	2779	2780	2781	2782	2783	2784	2785	2786	2787	2788	2789	2790	2791	2792	2793	2794	2795	2796	2797	2798	2799	2800	2801	2802	2803	2804	2805	2806	2807	2808	2809	2810	2811	2812	2813	2814	2815	2816	2817	2818	2819	2820	2821	2822	2823	2824	2825	2826	2827	2828	2829	2830	2831	2832	2833	2834	2835	2836	2837	2838	2839	2840	2841	2842	2843	2844	2845	2846	2847	2848	2849	2850	2851	2852	2853	2854	2855	2856	2857	2858	2859	2860	2861	2862	2863	2864	2865	2866	2867	2868	2869	2870	2871	2872	2873	2874	2875	2876	2877	2878	2879	2880	2881	2882	2883	2884	2885	2886	2887	2888	2889	2890	2891	2892	2893	2894	2895	2896	2897	2898	2899	2900	2901	2902	2903	2904	2905	2906	2907	2908	2909	2910	2911	2912	2913	2914	2915	2916	2917	2918	2919	2920	2921	2922	2923	2924	2925	2926	2927	2928	2929	2930	2931	2932	2933	2934	2935	2936	2937	2938	2939	2940	2941	2942	2943	2944	2945	2946	2947	2948	2949	2950	2951	2952	2953	2954	2955	2956	2957	2958	2959	2960	2961	2962	2963	2964	2965	2966	2967	2968	2969	2970	2971	2972	2973	2974	2975	2976	2977	2978	2979	2980	2981	2982	2983	2984	2985	2986	2987	2988	2989	2990	2991	2992	2993	2994	2995	2996	2997	2998	2999	3000	3001	3002	3003	3004	3005	3006	3007	3008	3009	3010	3011	3012	3013	3014	3015	3016	3017	3018	3019	3020	3021	3022	3023	3024	3025	3026	3027	3028	3029	3030	3031	3032	3033	3034	3035	3036	3037	3038	3039	3040	3041	3042	3043	3044	3045	3046	3047	3048	3049	3050	3051	3052	3053	3054	3055	3056	3057	3058	3059	3060	3061	3062	3063	3064	3065	3066	3067	3068	3069	3070	3071	3072	3073	3074	3075	3076	3077	3078	3079	3080	3081	3082	3083	3084	3085	3086	3087	3088	3089	3090	3091	3092	3093	3094	3095	3096	3097	3098	3099	3100	3101	3102	3103	3104	3105	3106	3107	3108	3109	3110	3111	3112	3113	3114	3115	3116	3117	3118	3119	3120	3121	3122	3123	3124	3125	3126	3127	3128	3129	3130	3131	3132	3133	3134	3135	3136	3137	3138	3139	3140	3141	3142	3143	3144	3145	3146	3147	3148	3149	3150	3151	3152	3153	3154	3155	3156	3157	3158	3159	3160	3161	3162	3163	3164	3165	3166	3167	3168	3169	3170	3171	3172	3173	3174	3175	3176	3177	3178	3179	3180	3181	3182	3183	3184	3185	3186	3187	3188	3189	3190	3191	3192	3193	3194	3195	3196	3197	3198	3199	3200	3201	3202	3203	3204	3205	3206	3207	3208	3209	3210	3211	3212	3213	3214	3215	3216	3217	3218	3219	3220	3221	3222	3223	3224	3225	3226	3227	3228	3229	3230	3231	3232	3233	3234	3235	3236	3237	3238	3239	3240	3241	3242	3243	3244	3245	3246	3247	3248	3249	3250	3251	3252	3253	3254	3255	3256	3257	3258	3259	3260	3261	3262	3263	3264	3265	3266	3267	3268	3269	3270	3271	3272	3273	3274	3275	3276	3277	3278	3279	3280	3281	3282	3283	3284	3285	3286	3287	3288	3289	3290	3291	3292	3293	3294	3295	3296	3297	3298	3299	3300	3301	3302	3303	3304	3305	3306	3307	3308	3309	3310	3311	3312	3313	3314	3315	3316	3317	3318	3319	3320	3321	3322	3323	3324	3325	3326	3327	3328	3329	3330	3331	3332	3333	3334	3335	3336	3337	3338	3339	3340	3341	3342	3343	3344	3345	3346	3347	3348	3349	3350	3351	3352	3353	3354	3355	3356	3357	3358	3359	3360	3361	3362	3363	3364	3365	3366	3367	3368	3369	3370	3371	3372	3373	3374	3375	3376	3377	3378	3379	3380	3381	3382	3383	3384	3385	3386	3387	3388	3389	3390	3391	3392	3393	3394	3395	3396	3397	3398	3399	3400	3401	3402	3403	3404	3405	3406	3407	3408	3409	3410	3411	3412	3413	3414	3415	3416	3417	3418	3419	3420	3421	3422	3423	3424	3425	3426	3427	3428	3429	3430	3431	3432	3433	3434	3435	3436	3437	3438	3439	3440	3441	3442	3443	3444	3445	3446	3447	3448	3449	3450	3451	3452	3453	3454	3455	3456	3457	3458	3459	3460	3461	3462	3463	3464	3465	3466	3467	3468	3469	3470	3471	3472	3473	3474	3475	3476	3477	3478	3479	3480	3481	3482	3483	3484	3485	3486	3487	3488	3489	3490	3491	3492	3493	3494	3495	3496	3497	3498	3499	3500	3501	3502	3503	3504	3505	3506	3507	3508	3509	3510	3511	3512	3513	3514	3515	3516	3517	3518	3519	3520	3521	3522	3523	3524	3525	3526	3527	3528	3529	3530	3531	3532	3533	3534	3535	3536	3537	3538	3539	3540	3541	3542	3543	3544	3545	3546	3547	3548	3549	3550	3551	3552	3553	3554	3555	3556	3557	3558	3559	3560	3561	3562	3563	3564	3565	3566	3567	3568	3569	3570	3571	3572	3573	3574	3575	3576	3577	3578	3579	3580	3581	3582	3583	3584	3585	3586	3587	3588	3589	3590	3591	3592	3593	3594	3595	3596	3597	3598	3599	3600	3601	3602	3603	3604	3605	3606	3607	3608	3609	3610	3611	3612	3613	3614	3615	3616	3617	3618	3619	3620	3621	3622	3623	3624	3625	3626	3627	3628	3629	3630	3631	3632	3633	3634	3635	3636	3637	3638	3639	3640	3641	3642	3643	3644	3645	3646	3647	3648	3649	3650	3651	3652	3653	3654	3655	3656	3657	3658	3659	3660	3661	3662	3663	3664	3665	3666	3667	3668	3669	3670	3671	3672	3673	3674	3675	3676	3677	3678	3679	3680	3681	3682	3683	3684	3685	3686	3687	3688	3689	3690	3691	3692	3693	3694	3695	3696	3697	3698	3699	3700	3701	3702	3703	3704	3705	3706	3707	3708	3709	3710	3711	3712	3713	3714	3715	3716	3717	3718	3719	3720	3721	3722	3723	3724	3725	3726	3727	3728	3729	3730	3731	3732	3733	3734	3735	3736	3737	3738	3739	3740	3741	3742	3743	3744	3745	3746	3747	3748	3749	3750	3751	3752	3753	3754	3755	3756	3757	3758	3759	3760	3761	3762	3763	3764	3765	3766	3767	3768	3769	3770	3771	3772	3773	3774	3775	3776	3777	3778	3779	3780	3781	3782	3783	3784	3785	3786	3787	3788	3789	3790	3791	3792	3793	3794	3795	3796	3797	3798	3799	3800	3801	3802	3803	3804	3805	3806	3807	3808	3809	3810	3811	3812	3813	3814	3815	3816	3817	3818	3819	3820	3821	3822	3823	3824	3825	3826	3827	3828	3829	3830	3831	3832	3833	3834	3835	3836	3837	3838	3839	3840	3841	3842	3843	3844	3845	3846	3847	3848	3849	3850	3851	3852	3853	3854	3855	3856	3857	3858	3859	3860	3861	3862	3863	3864	3865	3866	3867	3868	3869	3870	3871	3872	3873	3874	3875	3876	3877	3878	3879	3880	3881	3882	3883	3884	3885	3886	3887	3888	3889	3890	3891	3892	3893	3894	3895	3896	3897	3898	3899	3900	3901	3902	3903	3904	3905	3906	3907	3908	3909	3910	3911	3912	3913	3914	3915	3916	3917	3918	3919	3920	3921	3922	3923	3924	3925	3926	3927	3928	3929	3930	3931	3932	3933	3934	3935	3936	3937	3938	3939	3940	3941	3942	3943	3944	3945	3946	3947	3948	3949	3950	3951	3952	3953	3954	3955	3956	3957	3958	3959	3960	3961	3962	3963	3964	3965	3966	3967	3968	3969	3970	3971	3972	3973	3974	3975	3976	3977	3978	3979	3980	3981	3982	3983	3984	3985	3986	3987	3988	3989	3990	3991	3992	3993	3994	39
---------------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	----

VICK & PICKWELL

other two each fitted a capillary over a fang . Frequently, the snake voluntarily expelled venom into the capillaries. Subsequent pressure applied along the venom glands and ducts (maxillary margin) usually extruded an additional quantity of venom (table I). The product from a number of specimens was pooled for various tests and analyses. Fresh Pelamis venom was invariably water-clear.

C. Venom Treatment.

Depending on the tests planned, the freshly obtained whole venom was either refrigerated, frozen, or lyophilized. Fresh, unrefrigerated venom was, in some cases, placed on one of our gel filtration columns, submitted to pH determination, or percent solids and specific gravity measurements. For whole venom fractionation, we employed Sephadex G75-40 in either a 2 X 300 cm glass column for maximum single pass resolution (9) or a 2 X 150 cm glass column for more rapid separations and use aboard ship. Gel filtration columns were calibrated for molecular weight determinations using primary standards supplied for this purpose in kit form (Pharmacia).

D. Test Animals.

All basic toxicity tests were performed on adult male and female albino mice (age 8 to 10 weeks) from the Walter Reed randomly bred ICR/FG strain. Final toxicity assays employed either male mice weighing 30 to 35 grams, or female mice weighing 25 to 30 grams. Male and female mice were never mixed in toxicity tests. The LD50 was estimated using the graphic method of Miller and Tainter (10) and the statistical procedures described by Finney (11).

Stock venom test solutions were made up in cold physiological saline. Dilutions from stock were made so that venom concentration was contained in a standard volume for injection equivalent to approximately 1 percent of mouse body weight. Graded doses of the venom were injected into a tail vein of the mice. For each venom dose tested, 10 mice were used. The LD50 evaluation was determined from several pooled assays performed over a 24-hour period. The mice were closely observed for symptomatology and death for 4 hours following injection. Deaths occurring within 48 hours of injection were included in the evaluation of the LD50.

Five lots of sea snake antivenin, batch numbers 549-009(1964) 011 1(1967), 020-1(1969) 23-1(1970) and 25-1(1972), obtained from the Commonwealth Serum Laboratories, Australia were tested for efficacy

VICK & PICKWELL

against sea snake venom in Walter Reed ICR strain mice (Table IV)

Additional pharmacological tests were performed on adult beagle dogs (Army test stock) weighing 10 ± 2 kg. In these animals, the venom was injected into a indwelling catheter in the femoral vein. A single primate test was run on a 6-kg male Rhesus monkey. In all these tests, the dogs and the monkey were anesthetized with sodium pentobarbital (30 mg/kg). Tests for vasoconstrictive and cardiotoxic effects of the venom were performed on the isolated perfused dog heart following the method of Vick and Herman⁽¹²⁾.

RESULTS

A. Venom Production.

Bahia Banderas appears to be the northermost site of a year-round population of the Pelamis in the eastern Pacific and in addition, it functions as a nursery area where gravid females congregate to give birth to their free-swimming, precocious young. Consequently, the majority of the specimens listed in table-I were gravid and, therefore, probably heavier than non-gravid specimens of comparable length.

Even when there was no evidence for a broken or unregenerated fang, the venom production from right and left venom glands tended to be unequal by as much as a factor of two or more (table I). Frequently, a varying amount of venom would be voluntarily injected into the capillaries by the snake accompanied by visible contractions of the marginal maxillary musculature. Following this venom expulsion, an additional amount was usually available by milking. Venom production from apparently healthy, vigorous adult snakes ranged from a few lambda to a maximum of 33 lambda, but averaged 15 to 20 lambda per snake (table I; lambda = microliter). The venom was always waterclear and colorless except for the final few lambda expelled by milking which were occasionally cloudy with cellular debris.

B. Whole Venom Toxicity.

Fresh, liquid Pelamis venom (batch 1, refrigerated at 5 degrees C for no more than 4 days from time of milking to time of testing) was found to have an LD50 of 0.092 mg/kg in adult male mice (table II) based on an estimated 15.3 percent solids.* A second pooling of Pelamis whole venom that had been lyophilized (batch 2) showed an LD50 of 0.111 mg/kg in adult female mice (table III). Until

VICK & PICKWELL

further testing we cannot say whether the apparent difference in toxicity of the two batches may be ascribed to a loss of activity in the freeze dried venom, a possible greater resistance on the part of female mice as has been shown for female over male mice in the Malayan pit viper venom or some other factor.

A limited series of toxicity tests on dogs in conjunction with pharmacological evaluation suggested a somewhat lower LD50 for whole Pelamis venom (fresh, liquid) in dogs than in mice, but insufficient data are available for valid statistical treatment. Using dogs, Vick (13) found the lowest LD50 (0.05 mg/kg) yet reported for any sea snake venom.

C. Toxicity of Major Venom Fractions.

Separation of Pelamis whole venom into its main components on our gel columns produced two major fractions (one a composite) together constituting about 90 percent of the total venom, and a series of minor components* (figure 1).

In the first of two series of tests on lyophilized Pelamis venom fractions, the cuts were obtained and labeled as indicated in figure 1. Although there was subsequent evidence of possible degradation in some of the fractions due to an error in handling during transport, both fractions 2 and 3 were found to be toxic for male albino mice (LD50 of 7.5 and 0.89 mg/kg, respectively). Neither fraction was as toxic as the fresh whole venom, obviously. Fraction 1 was found to be totally nontoxic for mice at concentrations from 50 to 500 mg/kg. Whereas fraction 3 from this batch proved quickly fatal for a single dog at the concentration tested, the same dose of fraction 2 in a second dog of equivalent size was not fatal.

In a second series of tests, cuts were made on the fractionated venom. In this case, some of the minor fraction residues were also tested. Again, only fractions 2 and 3 demonstrated toxic activity, whereas the remaining components were nontoxic at the maximum concentrations tested. Careful assessment of the LD50's for fractions 2 (0.201 mg/kg) and 3 (0.086 mg/kg) in female albino mice indicated that fraction 3 was more toxic than whole Pelamis venom. Fraction 2 is about as toxic as whole venom from the common cobra (Naja naja) (14).

D. Pharmacology.

VICK & PICKWELL

1. Whole Venom.

The symptoms of envenomation in male and female mice were essentially identical. The dose-response relationship was very steep, typical of highly toxic substances. As the venom dose decreased survival times increased. Major symptoms observed included deep and rapid respiration, always accompanied by severe mydriasis. At doses greater than the LD50 level, the mice were lethargic or semi-comatose and died almost instantaneously. At doses near the LD50, the symptoms of labored and rapid respiration occurred approximately 15 minutes postinjection, accompanied by mydrdriasis. The heart beat seemed to increase. The mice at this stage appeared severely depressed and lethargic, showing impaired reflexes and occasional convulsions. As the effects progressed, respiration decreased and became more shallow until death occurred. If the mice remained in this state for more than 2 hours following injection, they usually recovered.

Physiological data obtained from dogs receiving varying doses of (13) Pelamis whole venom (batch 1) paralleled results obtained by Vick with dogs receiving venom of the sea snakes *Laticauda laticaudata* and *L. semifasciata*. This was particularly apparent for the arterial blood pressure, heart rate, and EKG, all of which remained unchanged for periods of minutes even at the highest venom doses. In common with envenomated mice, the respiration rate in some test dogs was unaffected at first, only gradually increasing in the final preterminal minutes as inspiratory volume declined. In the dogs receiving the highest doses, however, respiration rate remained almost unchanged until time of death.

In animals nearing death, arterial pressure rose simultaneously with a decrease in inspiratory volume while heart rate gradually declined. The abruptness of onset of these latter events and the speed with which they developed seemed directly related to venom dosage. The EKG wave form remained almost unchanged until time of termination.

Physiological events in poisoned monkeys were practically the same as in dogs except that bradycardia developed very abruptly and only in the terminal phase.

A series of graded doses of Pelamis whole venom (batch 1) was tested on the isolated, perfused dog heart. At dose levels equivalent to 0.5, 1.0, 1.5, 2.0, and 2.5 mg/kg of intact animal, no change in rate or force of contraction was shown by the perfused, beating

VICK & PICKWELL

hearts.

The effectiveness of sea snake antivenin against the venom of 3 sea snake species is shown in Table IV. Results indicate that the antivenin is highly effective against the venoms of *Laticauda laticaudata*, *Laticauda semifasciata*, and *Pelamis platurus* and in addition retains its potency for at least 8 years.

2. Venom Fractions.

Although there were difference in toxicity between fractions 2 and 3 of *Pelamis* venom, and between these fractions and whole venom, the symptoms produced in mice by each fraction were the same as those elicited by whole venom. That is, the mice envenomated with either fraction in the toxic range showed dyspnea, mydriasis, and impairment of reflexes, particularly the righting reflex. Death, with whole venom, appeared to occur from respiratory failure.

In the dog poisoned with fraction 3, the sequence of events was nearly identical to the sequence seen with high doses of whole venom. In the animals receiving fraction 2, no symptoms directly attributable to the effects of envenomation were observed throughout the 3-hour experiment.

DISCUSSION.

A. Cause of Death from *Pelamis* Venom.

From the foregoing, it appears that *Pelamis* venom acts to produce respiratory arrest in animals critically poisoned. No evidence for cardio toxicity was found in whole venom or its toxic fractions. This is essentially in agreement with earlier work indicating respiratory arrest to be a major cause of death in experimental animals^(15,16) and in human beings who are victims of sea snake bite. Respiratory failure arising from myoneural junction blockade at the level of the diaphragm was the cause of death in experimental animals injected with whole venom from the sea snakes *Laticauda laticaudata* and *L. semifasciata*⁽¹³⁾.

B. Hazard to Man from Bites of *Pelamis*.

The venom of *Pelamis* is somewhat less toxic than that of other sea snakes studied in recent years^(15,16). In some cases, its

VICK & PICKWELL

potency is equal to or greater than that of other sea snake species (8,16,18,20). However, few toxicity studies of sea snake venom published to date have been done with optimal numbers of laboratory animals, particularly mice, from suitably homogeneous populations; or perhaps this important facet has been left undisclosed by the respective authors (exceptions -Vick (15) and Tu and Ganthavorn) (20). Data obtained from test groups of fewer than 30 are frequently not statistically reliable, (21,22) whereas populations of widely divergent size or age, or mixed gender* can all contribute to substantial deviations in LD50 values. Nonetheless, keeping in mind these difficulties, it appears that Pelamis venom toxicities are equal to or greater than those determined by others for this species.

Pelamis produces much less venom than does other species of sea snakes. Even in large specimens (table 1) the quantity available for injection will not be fatal to an adult man if the mouse toxicity data can be extrapolated to mammals, including man. Tests on monkeys suggest that primates may be less susceptible to sea snake venom than are mice. This becomes particularly true considering the work of Reid (16) who was of the opinion that sea snakes infrequently inject venom when biting in defense, and also considering the further probability that a sea snake is unlikely to deliver its total venom supply in a single bite. Under certain circumstances, however, it is possible to visualize how a bite from a large Pelamis might prove serious for a small adult, especially an infirm or elderly person, or a child.

C. Aggressive Behavior of Pelamis Toward Man.

Reid (16) and Barme (19) have documented the incidence of sea snake bites among native fishermen of Southeast Asia. The bites occurred most frequently around the fishermen's hands and arms while they were handling nets and sorting fish, or on their feet and ankles while they were wading in their nets, stepping on or otherwise disturbing unseen snakes in the muddy water. Pelamis was not implicated as the source of envenomation in these reports.

Pickwell (23) has summarized the available information concerning the behavior of sea snakes toward swimmers and divers. The evidence indicates that swimmers on the surface or divers in transit are unmolested by the sea snakes they encounter although a number of instances have been reported of swimmers being followed some distance by sea snakes. Divers working in static situations such as salvage operations have, however, been bitten by sea snakes. There are no verified reports of loss of life under these circumstances, and it is

VICK & PICKWELL

not clear what attracts the snakes to such relatively large animals as the stationary human divers.

Neither has Pelamis been implicated in any of the diving and swimming encounters reported to us. In contrast to the behavior of other species, Pelamis has not been reported to approach or to bite man. This may be largely because of its life habits and habitat (24) which include spending much of its life at the sea surface where it alone among the 50 species of sea snakes is able to feed on small pelagic fishes. Pelamis thus does not dive for feeding or hunting purposes and is not tied to inshore areas where wading fishermen or divers would most frequently be encountered. Nevertheless, Pelamis does occur in such areas as well as on the open sea, and we have had opportunity to test its possible aggressiveness or curiosity for human divers and swimmers.

In Bahia Banderas, Mexico we have swum with Pelamis where they occurred in some numbers several hundred yards offshore. None of us were ever approached by the sea snakes even when we had been treading water in a stationary position for some time. While in the water, one of our group once attempted to photograph specimens of this species. In two separate instances the snakes swam away from the diver just rapidly enough to maintain their distance, regardless of how fast the diver swam in pursuit. It seems, on the basis of these experiences, the Pelamis is not among the sea snake species likely to be aggressive or, for that matter, annoyingly curious about people.

There are very few accounts of human deaths from Pelamis envenomation. Such accounts as do exist are impossible to verify and tend to be anecdotal. An example of these is Beck's (25) account of a young diver (working in the Torres Straits, Cape York Peninsula, Australia) who was bitten on the finger by a yellow-bellied sea snake (presumably Pelamis). The man became convulsive and died 48 hours later despite the fact the finger was amputated within an hour of receiving the bite.

Halstead (26) in his monumental treatise on venomous and poisonous marine animals cited a total of four deaths from the bite of Pelamis. He also included more generalized statements from the literature regarding the mortal fear in which this snake is held in such areas as Taiwan. Swaroop and Grab (27) in their summary on worldwide snake bite mortality stated that deaths from Pelamis bite have occurred near Mozambique and along the coast of Central America. No data or references were given, however, and we do not know the

VICK & PICKWELL

source of these statements. Reid and Lim (28) in the course of a survey of Malay fishing villages, learned of the "ular belerang," a mythical sea snake feared more than any other by the natives. The description of this snake (claimed to have been seen by a few of the native people) suggests a young Pelamis (belerang is Malay for sulfur). Fayrer (29) quoted a Mr. Stewart as saying that Indian natives in the region of Puri, in the state of Orissa, on the east coast, believe Pelamis to be the most deadly of all sea snakes. As "kullunder samp," this species figures prominently in local folklore.

d. Pelamis on the Beach.

Although it is able to feed and survive in the open sea and is not committed to shallow water, Pelamis not only occurs near shore in such places as Bahia Banderas, Mexico, and the Bay of Panama but frequently is washed ashore in these areas (24,30-32). Once on the beach, the sea snake is comparatively helpless since it lacks the broad ventral scutes possessed by land snakes. We have reports of large numbers of Pelamis stranded along the beaches of Panama and northern Peru, and we have observed and counted stranded Pelamis along the beaches at Puerto Vallarta, Bahia Banderas (24). Many of these snakes were still vigorous in the early morning when we made our counts, and they readily swam away when returned to the water. The possibility of receiving a bite from accidentally treading upon a beached Pelamis seemed real enough since the beaches in this resort area are heavily patronized by Mexican and American tourists alike. Yet our conversations with the town mayor, a local physician, and the port captain for Puerto Vallarta failed to disclose any incidents of sea snake bite within their memory. Evidently, the sea snake fails to intimidate the tourists frequenting the beaches of this popular resort and, at least in this area, has caused no fatalities or serious cases of envenomation.

e. Sea Snake Antivenin.

The currently available antivenin is effective against at least 3 species of sea snake and most probably many more. In addition it retains its potency for at least 8 years which makes it quite useful in actual field situations. Early results in our laboratory further indicate that short term artificial respiration, in addition to antivenin therapy, may increase survivor populations.

VICK & PICKWELL

LITERATURE CITED

1. Cropp B. Sea Snakes. Oceans Magazine 3(2), 48-54 (1970).
2. Dunson, W. A. The Sea Snakes are Coming. Natural History Magazine 80(9), 52-61 (1971).
3. MacLeish K. Diving with Sea Snakes. National Geographic 141(4), 565-578 (1972).
4. Weathersbee, C. Linking the Oceans. Science News 94(23), 578-581 (1968).
5. Rubinoff I., and Kropach, C. Differential Reactions of Atlantic and Pacific Predators to Sea Snakes. Nature, London 228, 1288-1290 (1970).
6. Gilluly, R. H. Consequences of a Sea-level Canal Science News 99(3) 52-53 (1971).
7. Bailey, D. A Thermal Barrier for Snakes. Biomed News 12-70. December 1970.
8. Barme M. Venomous Sea Snakes (Hydrophiidae) in Venomous Animals and Their Venoms. Vol. 1. Venomous Vertebrates. W. Bucherl E. E. Buckley, and V. Deulofeu, eds., pp 285-308. Academic Press, New York, New York. 1968.
9. Shipman, W. H., and Cole, L. J. A Surfactant Bee Venom Fraction: Separation on a Newly Devised Constant Flow Rate Chromatographic Column and Detection by Changes in Effluent Drop Volume. Anal. Biochem. 29, 490-497 (1969).
10. Miller, L. C. and Tainter M. L. Estimation of the ED50 and Its Error by Means of Logarithmic-Probit Graph Paper. Proc. Soc. Exp. Biol. Med. 57, 261-264 (1944).
11. Finney, D. J. Probit Analysis. 3d Ed. 333 pp. Cambridge University Press. 1971.
12. Vick, J. A., and Herman, E. H. An Isolated Dog or Monkey Heart Preparation for Studying Cardioactive Compounds. Pharmacology 6, 290-299 (1971).

VICK & PICKWELL

13. Vick, J. A. Venomous Sea Snakes. pp 974-975 in Poisonous and Venous Marine Animals of the World. Vol. 3 - Vertebrates Continued. B. W. Halstead. US Government Printing Office, Washington, DC. 1970.

14. Vick, J. A., and Grenan, M. M. Evaluation of Antivenoms for Common Cobra (*Naja naja*) and Blue Krait (*Bungarus caeruleus*) Venoms. In Handbook of Dangerous Animals for Field Personnel, G. V. Pickwell and W. E. Evans, eds., Naval Undersea Center, San Diego. California Technical Report. 1972.

15. Rogers. L. On the Physiological Action of the Poison of the Hydrophiadae. Part II. Action on the Circulatory, Respiratory, and Nervous Systems. Royal Society, London. Proceedings 72, 305-319 (1903).

16. Reid, H. A. Sea-Snake Bite Research. Trans Roy. Soc. Trop. Med. Hyg. 50(6), 517-542 (1956).

17. Reid H. A. Snake Bite in the Tropics. British Med. J. 3, 359-362 (1968).

18. Carey, J. E., and Wright, E. A. The Toxicity and Immunological Properties of Some Sea-Snake Venoms with Particular Reference to That of *Enhydrina schistosa*. Trans. Roy. Soc. Trop. Med. Hyg. 54(1), 50-67 (1960).

19. Barme, M. Venomous Sea Snakes of Viet Nam and Their Venoms. pp 373-378 in Venomous and Poisonous Animals and Noxious Plants of the Pacific Region H. L. Keegan and W. V. Macfarlane, eds., Pergamon (Macmillan), New York, New York. 1963.

20. Tu, A. T. and Ganthavorn, S. Immunological Properties and Neutralization of Sea Snake Venoms from Southeast Asia. Am. J. Trop. Med. Hyg. 18(1) 151-154 (1969).

21. Trevan, J. W. The Error of Determination of Toxicity. Royal Society, London. Proceedings. B. 101(712), 483-514 (1927).

22. Russell F. E. To be or Not to be... the LD50. Editorial. Toxicon. 4, 81-83 (196).

23. Pickwell, G. V. Sea Snakes of Viet Nam and Southeast Asia. In Handbook of Dangerous Animals for Field Personnel G. V. Pickwell and W. E. Evans, eds., Naval Undersea Center, San Diego,

VICK & PICKWELL

California Technical Report. 1972.

24. Pickwell, G. V. The Venomous Sea Snakes. Fauna. The Zoological Magazine. No. 4, pp 17-32 1972.

25. Becke, L. Neath Austral Skies. pp 305-310. John Milne, London 1909.

26. Halstead, B. W. Poisonous and Venomous Marine Animals of the World. Vol. 3. pp 627-681 - Vertebrates continued US Government Printing Office, Washington, DC. 1970.

27. Swaroop, S., and Grab, B. Snakebite Mortality in the World. Bull. World Health Org. 10, 35-76 (1954).

28. Reid, H. A., and Lim, K. J. Sea-Snake Bite. A Survey of Fishing Villages in Northwest Malaya. British Med. J. ii, 1266-1272 (1957).

29. Fayrer, J. the Thanatophidia of India. pp 131,133. J. & A. Churchill, London, 1872.

30. Viquez, C. Distribucion geografica de nuestras serpientes venenosas. Inst. Defensa Cafe Costa Rica. Revista. II(87), 608-611 (1942).

31. Dunson W. A., and Ehlert, G. W. Effects of Temperature, Salinity, and Surface Water Flow on Distribution of the Sea Snake Pelamis. Limnol. Oceanog. 16. 845-853 (1971).

32. Kropach, C. Pelamis platurus as a Potential Colonizer of the Caribbean Sea. Bull. Biol. Soc. Wash. 2, 267-269 (1972)

33. Rogers, L. On the Physiological Action of the Poison of the Hydrophidae. Royal Society, London Proceedings 71, 481-496 (1902 1903).

34. Nauck E. G. Untersuchungen uber das Gift einer Seeschlange (Hydrus platurus) des Pazifischen Ozeans Deutsche Tropenmed. Zeit. 33, 167-170 (1929).

Table I Venom Production by Newly Captured Adult, Female Sea Snakes (*Pelamis platurus*)^a

Specimen number	Length ^b	Weight ^b	Venom produced ^c				Total	
			Left fang		Right fang		Liquid venom	Equivalent dry venom ^d
			Voluntary	Milked	Voluntary	Milked		
	cm	gm	lambda				lambda	mg
1	70	160	0	5.8	4.2	1.5	12	1.8
2	75	180	9.6	1.5	5.8	0	17	2.6
3	83	310	8.1	7.7	15.8	1.5	33	5.0
4	75	200	12.7	3.8	13.1	0	30	4.6
5	68	150	1.1	3.8	0	11.1	16	2.4
6	71	160	1.9	1.9	1.9	0.4	6	0.9
7	72	170	4.2	0.8	0.8	2.3	8	1.2
8	80	200	10.0	2.3	4.6	2.7	20	3.1
9	70	160	3.8	2.7	5.8	0	12	1.8
10	77	210	14.6	0.8	6.9	0	22	3.4
11	75	210	5.8	0.8	0.4	3.1	10	1.5
12	73	190	13.1	1.1	8.8	5.0	28	4.3
13	62	110	4.2	2.7	3.8	5.8	17	2.6

^a Data obtained aboard R/V CAPE on Bahia Banderas, Mexico, 21 to 24 March 1971.^b Lengths, ± 1 cm, approx; weights, ± 10 gm.^c 1 lambda = 1 microliter.^d Based on 15.3 average percent solids in liquid *Pelamis* venom (Shipman and Pickwell, in preparation).

Table II LD50* Assay - *Pelamis* Liquid Whole Venom in Mice**

Dose group	Dosage	Number of mice	Number dead	Mortality	Mean survival time
	mg dry venom/kg			%	min
1	0.20	5	5	100	27
2	0.12	10	7	70	40
3	0.10	10	6	70	68
4	0.095	10	6	60	68
5	0.086	10	3	30	51
6	0.078	10	2	20	63

*This was fresh, liquid whole venom, batch 1 (see text). Mice were all males, 8 weeks old, 30 to 35 gm of the Walter Reed ICR strain.

**LD50 = 0.092 mg/kg.

Table III LD50* Assay - *Pelamis* Lyophilized Whole Venom in Mice**

Dose group	Dosage	Number of mice	Number dead	Mortality	Mean survival time
	mg dry venom/kg			%	min
1	0.5	5	5	100	5
2	0.2	5	5	100	23
3	0.14	10	10	100	33
4	0.13	30	25	83	40
5	0.12	30	19	63	48
6	0.11	20	11	55	64
7	1.10	20	5	25	64
8	0.075	10	0	-	-

*This was *Pelamis* venom batch 2 (see text). Mice were all females, 9 to 10 weeks old, 25 to 30 gm, of the Walter Reed ICR strain.

**LD50 = 0.111 mg/kg.

Table IV Effectiveness of Sea Snake Antivenin* Against the Venom of Three Heterologous Sea Snake Species

Species	Date of test	Venom LD50 for mice	Venom LD99 for mice	Antivenin and venom LD50	Potency factor**	Venom (mg) 1 ml antivenin	Neutralized by 1 unit antivenin	Antivenin		Strength units/ml
								Batch No.	Date of batch	
<i>Pelamis platurus</i>	September 1972	0.11	0.15	1.06	9.6	0.1	0.004	25-1	February 1972	26.6
								23-1	June 1970	27.5
								020-1	March 1969	23.1
								011-1	26 June 1967	44.8
								549-009	24 November 1964	34.4
<i>Laticauda laticaudata</i>	December 1972 January 1973	0.16	0.26	1.31	8.2	0.3	0.011	25-1	February 1972	26.6
								23-1	June 1970	27.5
								020-1	March 1969	23.1
								011-1	26 June 1967	44.8
								549-009	24 November 1964	34.4
<i>Laticauda semifasciata</i>	April 1973 May 1973	0.30	0.48	1.90	6.3	0.4	0.015	25-1	February 1972	26.6
								23-1	June 1970	27.5
<i>Laticauda semifasciata</i>				1.65	5.5	0.4	0.017	020-1	March 1969	23.1
								011-1	26 June 1967	44.8
								549-009	24 November 1964	34.4

* Commonwealth Serum Laboratories, Australia - equine, liquid, sea snake antivenin (AV) produced from the venom of the common sea snake, *Enhydrius schistosus*, five batches of varying age. All injections IV in mice.

** Potency factor = $\frac{\text{LD50 AV} + \text{venom}}{\text{LD50 venom only}}$

† One unit of Commonwealth Serum Laboratories' sea snake antivenin is defined as that quantity which will neutralize 0.01 mg of *Enhydrius schistosus* venom.




Figure 1 Separation of *Pelamis* Venom Components by Single Pass Gel Filtration on a 2 x 150-cm Column